

Interpolation Formulas for the $\eta\gamma$ and $\eta'\gamma$ Transition Form Factors

Th. Feldmann* and P. Kroll

Fachbereich Physik, Universität Wuppertal, D-42097 Wuppertal, Germany
(to be published in *Physical Review D*)

The new CLEO and LEP data [1,2] on the $\eta\gamma$ and $\eta'\gamma$ transition form factors have renewed the interest in simple interpolation formulas, valid at any value of momentum transfer. We are going to show that recent theoretical and phenomenological results on η - η' mixing [3–5] lead to two-pole forms, where each pole term resembles the Brodsky/Lepage interpolation formula for the $\pi\gamma$ case [6] and depends on the mixing and decay parameters in a simple fashion. The parameters, entering the occasionally used one-pole formulas, on the other hand, cannot be interpreted theoretically in a simple way.

Copyright 1998 by The American Physical Society.

I. INTRODUCTION

In 1995 the CLEO collaboration has presented their preliminary data on pseudoscalar meson-photon transition form factors (i.e. on the processes $\gamma\gamma^* \rightarrow P$, $P = \pi, \eta, \eta', \dots$) at large space-like momentum transfer, Q^2 , for the first time [7]. One of the exciting aspects of these form factors is that they possess well-established asymptotic limits [8,9], which are related to mesonic decay constants only and behave as $1/Q^2$. On the other hand, at vanishing momentum transfer, $Q^2 = 0$, the values of the form factors are fixed by PCAC and the chiral anomaly and appear to be proportional to the inverse of the mesonic decay constants. The investigation of the transition form factors between the two limiting cases provides important information on the wave functions of the pseudoscalar mesons.

In case of the π^0 , where the decay constant is known from the leptonic π decay, Brodsky and Lepage (BL) [6] proposed a simple parameter-free formula which interpolates between the PCAC value at $Q^2 = 0$ and the QCD prediction for $Q^2 \rightarrow \infty$

$$F_{P\gamma}^{BL}(Q^2) = \frac{6 C_P f_P}{Q^2 + 4\pi^2 f_P^2}, \quad (P = \pi) \quad (1.1)$$

($C_\pi = 1/(3\sqrt{2})$ represents a charge factor). This formula works fairly well although it is not perfect, and may be used as an estimate of the $\pi\gamma$ transition form factor. Since the x -dependence (where x is the usual momentum fraction the quark carries) of the pion's wave function seems to be close to that of the asymptotic form [10,11], the phenomenological success of the BL interpolation formula is not a surprise. Note that the vector meson dominance model also provides a formula like (1.1). It is very close to the BL interpolation formula numerically since M_ρ approximately equals $2\pi f_\pi$. Even the pole residue agrees rather well with $6 C_\pi f_\pi$ numerically.

In case of the $\eta\gamma$ and $\eta'\gamma$ the situation is more complicated due to the η - η' mixing which has to be taken into account properly. η - η' mixing has been the subject of considerable interest and being examined in many investigations, e.g. [12–14] and references therein. Recent investigations [3–5] have shown, that the decay constants in the η - η' system are not adequately described in the octet-singlet scheme with one and the same angle for the particle state mixing and the parameterization of the decay constants. As has become clear very recently [5], a mixing-scheme where one takes as the basis the states according to their quark flavor decomposition is better suited for the description of the η and η' decay constants and meets all theoretical and phenomenological requirements. It also allows for proper definitions of the desired generalizations of (1.1) to the η and η' cases. We define the quark basis by the two states η_q and η_s

*Supported by *Deutsche Forschungsgemeinschaft*

composed of the valence quarks $q\bar{q} = (u\bar{u} + d\bar{d})/\sqrt{2}$ and $s\bar{s}$ (and corresponding higher Fock states), respectively. These states are related to the physical states by the transformation

$$\begin{pmatrix} \eta \\ \eta' \end{pmatrix} = U(\phi) \begin{pmatrix} \eta_q \\ \eta_s \end{pmatrix}, \quad U(\phi) = \begin{pmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{pmatrix}. \quad (1.2)$$

The four decay constants of the η and η' mesons, which are defined by

$$\langle 0 | J_{\mu 5}^i | P \rangle = i f_P^i p_\mu, \quad (i, j = q, s, P = \eta, \eta'), \quad (1.3)$$

can be expressed in terms of three parameters: the mixing angle ϕ and two decay constants f_q and f_s ,

$$\begin{pmatrix} f_\eta^q & f_\eta^s \\ f_{\eta'}^q & f_{\eta'}^s \end{pmatrix} = U(\phi) \begin{pmatrix} f_q & 0 \\ 0 & f_s \end{pmatrix}. \quad (1.4)$$

Here, the weak axial-vector currents are defined by $J_{\mu 5}^q = (\bar{u}\gamma_\mu\gamma_5 u + \bar{d}\gamma_\mu\gamma_5 d)/\sqrt{2}$ and $J_{\mu 5}^s = \bar{s}\gamma_\mu\gamma_5 s$. We repeat that in the conventional octet-singlet scheme the decay constants do NOT follow the pattern of state mixing, i.e. the angles appearing in the state mixing and in the parameterization of the decay constants are not identical [3–5,15]. On exploiting the divergences of the axial-vector currents, the basic mixing parameters ϕ , f_q and f_s are fixed to first order in the flavor symmetry breaking. In order to take into account higher order corrections the mixing parameters can be determined phenomenologically. In [5] the following values have been shown to give a consistent description of the available experimental data and to be in line with the constraints from the axial vector anomaly

$$\phi = 39.3^\circ \pm 1.0^\circ, \quad f_q = (1.07 \pm 0.02) f_\pi, \quad f_s = (1.34 \pm 0.06) f_\pi. \quad (1.5)$$

These phenomenological values agree within 10% with the theoretical values [5] and with the results from chiral perturbation theory [3]. We will use the values given in (1.5) for the following discussion of the $\eta\gamma$ and $\eta'\gamma$ transition form factors. In [5] it has already been found that they give a perfect description of the experimental $\eta\gamma$ and $\eta'\gamma$ form factor data above $Q^2 = 1 \text{ GeV}^2$ if the form factors are calculated within the modified hard-scattering approach [4] and if wave functions similar to the pion one are used.

II. GENERALIZED BL INTERPOLATION FORMULAS

Within the $q\bar{q}$ – $s\bar{s}$ mixing scheme the following generalization of (1.1) seems to be most natural ($P = \eta, \eta'$)

$$F_{P\gamma}^{BL}(Q^2) = \frac{6 C_q f_P^q}{Q^2 + 4\pi^2 f_q^2} + \frac{6 C_s f_P^s}{Q^2 + 4\pi^2 f_s^2}, \quad (2.1)$$

where the charge factors are $C_q = 5/9\sqrt{2}$, $C_s = 1/9$. At $Q^2 = 0$ it reproduces the PCAC formula (see e.g. [5]) which enters the two-photon widths of the η and η' mesons. For $Q^2 \rightarrow \infty$ QCD predicts the value of the form factors in terms of the decay constants f_P^i defined by (1.3). The latter represent wave functions of the lowest parton Fock states, $q\bar{q}$ and $s\bar{s}$, contributing to the η and η' mesons, taken at zero spatial separation, which appear in the perturbative calculation of the form factor (see e.g. [4]). The numerical factors in (2.1) are chosen in such a way that the asymptotic behavior, as predicted by QCD, is reproduced.

Comparison with experiment (see Fig. 1) reveals that the interpolation formulas (1.1) and (2.1) do not work badly. However, they should still be used with care: For momentum transfer of the order of a few GeV^2 the predicted curvatures of the form factors are stronger than the present data exhibit. On the other hand, the predictions obtained within the modified hard scattering approach are in perfect agreement with experiment [4,5,10] above 1 GeV^2 .

Our generalization of the BL interpolation formula (2.1) parallels the vector meson dominance model. This is so because the ω meson is nearly a pure $q\bar{q}$ state and mass degenerated with the ρ , while the ϕ meson is almost a pure $s\bar{s}$ state. In the vector meson dominance ansatz

$$F_{P\gamma}^{VMD}(Q^2) = \sum_{V=\rho,\omega,\phi} \frac{g_{PV\gamma} M_V^2}{f_V} \frac{1}{Q^2 + M_V^2} \quad (2.2)$$

one can effectively combine the first two terms into one because $M_\rho \simeq M_\omega$. This term can then be identified with the first term in (2.1), since $M_{\rho,\omega} \simeq 2\pi f_q$. The remaining contribution of the ϕ meson corresponds to the second term in (2.1) since $M_\phi \simeq 2\pi f_s$. Moreover, as a detailed analysis reveals, the corresponding residues in the generalized BL

interpolation formula and in the vector meson dominance ansatz approximately agree as well (for values of $g_{PV\gamma}$ and f_V see [16,17]).

Occasionally, one-pole BL interpolation formulas, like (1.1), are used for the η and η' cases [1,2,18,19] (see also the tables of particle data [20]). The corresponding parameters, C_P and f_P ($P = \eta, \eta'$), however, have no simple theoretical interpretation. In particular f_P , misleadingly called ‘decay constant’, does not represent a vacuum-particle matrix element of an appropriate axial-vector current and is, therefore, not related to basic decay properties of the η and η' mesons in a simple way. The requirement that the one-pole formulas interpolate between the PCAC results at $Q^2 = 0$ and the asymptotic perturbative QCD predictions, relates the parameters C_P and f_P to the decay constants f_P^i . Using the $q\bar{q}$ - $s\bar{s}$ mixing scheme one finds

$$\begin{aligned} C_P &= \sqrt{(C_q f_P^q + C_s f_P^s)(C_q f_P^q/f_q^2 + C_s f_P^s/f_s^2)} , \\ f_P &= \sqrt{\frac{C_q f_P^q + C_s f_P^s}{C_q f_P^q/f_q^2 + C_s f_P^s/f_s^2}} \end{aligned} \quad (2.3)$$

Despite of the fact that the parameters of the one-pole formula have a much more complicated interpretation than those of (2.1), the one-pole formula is of similar quality. Using, for instance, the values (1.5) for the basic mixing parameters to estimate C_P and f_P , one obtains results from the one-pole formula which are almost indistinguishable from what is found from (2.1).

Amettler et al. [21] also proposed a one-pole BL interpolation formula for the $\eta\gamma$ and $\eta'\gamma$ transition form factors. Since they implicitly assumed $C_P = C_\pi$, their formula only matches the PCAC result at $Q^2 = 0$ while it fails asymptotically and, hence, at large Q^2 .

III. SUMMARY

We have proposed BL interpolation formulas for the $\eta\gamma$ and $\eta'\gamma$ transition form factors which resemble the BL formula for the $\pi\gamma$ case. The formulas (2.1) are derived within the new mixing scheme advocated for in [5], where the η and η' mesons are expressed as linear combinations of orthogonal states η_q and η_s . Only in this mixing scheme the decay constants f_P^i ($i = q, s$, $P = \eta, \eta'$) can be described in terms of two basic decay constants f_q, f_s and the state mixing angle ϕ . This is a necessary requirement for the BL interpolation formula to be of the desired simple form. It is then rather similar to the vector meson dominance model, since the vector mesons show nearly ideal mixing, i.e. they distinguish between the $s\bar{s}$ and $q\bar{q}$ components in a natural way. Thus, a two-pole ansatz for the BL interpolation formula appears to be very natural. Its other advantage is that the basic mixing parameters, f_q, f_s, ϕ , enter in a simple fashion. With the values (1.5) for the basic mixing parameters the interpolation formulas (2.1) are in rough agreement with the experimental data [1,2,18,19,22]. Above $Q^2 = 1 \text{ GeV}^2$ the data are much better reproduced within a perturbative approach [4,5]. Despite of this, the interpolation formula is illustrative and may be used for quick estimates of the form factors.

It is also possible to specify one-pole BL interpolation formulas for the $\eta\gamma$ and $\eta'\gamma$ transition form factors. However, the parameters appearing in the one-pole formulas, have no simple theoretical interpretation; they depend on the basic mixing parameters in a very complicated way. In particular, the parameters f_P cannot be interpreted as ‘decay constants’ as is done occasionally.

-
- [1] CLEO collaboration, J. Gronberg *et al.*, Phys. Rev. **D57**, 33 (1998).
 - [2] L3 collaboration, M. Acciarri *et al.*, (1997), preprint CERN-PPE/97-110, submitted to *Phys. Lett. B*.
 - [3] H. Leutwyler, (1997), hep-ph/9709408, and R. Kaiser, diploma thesis, Bern 1997.
 - [4] T. Feldmann and P. Kroll, (1997), hep-ph/9711231, to be published in *Eur. Phys. J. C*.
 - [5] T. Feldmann, P. Kroll, and B. Stech, (1998), hep-ph/9802409.
 - [6] S. J. Brodsky and G. P. Lepage, Phys. Rev. **D24**, 1808 (1981).
 - [7] CLEO collaboration, V. Savinov *et al.*, (1995), in proceedings of the PHOTON 95 workshop, Sheffield (1995), eds. D.J. Miller *et al.*, World Scientific.
 - [8] T. F. Walsh and P. Zerwas, Nucl. Phys. **B41**, 551 (1972).
 - [9] G. P. Lepage and S. J. Brodsky, Phys. Rev. **D22**, 2157 (1980).

- [10] P. Kroll and M. Raulfs, Phys. Lett. **B387**, 848 (1996).
- [11] I. V. Musatov and A. V. Radyushkin, Phys. Rev. **D56**, 2713 (1997).
- [12] H. Fritzsch and J. D. Jackson, Phys. Lett. **66B**, 365 (1977).
- [13] N. Isgur, Phys. Rev. **D13**, 122 (1976).
- [14] F. J. Gilman and R. Kauffman, Phys. Rev. **D36**, 2761 (1987).
- [15] A. V. Kiselev and V. A. Petrov, Z. Phys. **C58**, 595 (1993).
- [16] O. Dumbrajs *et al.*, Nucl. Phys. **B216**, 277 (1983).
- [17] P. Ball, J. M. Frere, and M. Tytgat, Phys. Lett. **B365**, 367 (1996).
- [18] CELLO collaboration, H. J. Behrend *et al.*, Z. Phys. **C49**, 401 (1991).
- [19] TPC/2 γ collaboration, H. Aihara *et al.*, Phys. Rev. Lett. **64**, 172 (1990).
- [20] Particle Data Group, R. M. Barnett *et al.*, Phys. Rev. **D54**, 1 (1996).
- [21] L. Ametller, J. Bijnens, A. Bramon, and F. Cornet, Phys. Rev. **D45**, 986 (1992).
- [22] PLUTO collaboration, C. Berger *et al.*, Phys. Lett. **142B**, 125 (1984).




FIG. 1. Results for the $\pi\gamma$, $\eta\gamma$ and $\eta'\gamma$ transition form factors calculated using the modified hard scattering approach with asymptotic wave functions, and with the interpolation formulas (1.1,2.1). The straight lines indicate the asymptotic limit. The parameter values are $f_\pi = 131$ MeV, $\phi_P = 39.3^\circ$, $f_q = 1.07f_\pi$, $f_s = 1.34f_\pi$. Data are taken from [1,2,18,19,22].